


# Lagrangian Turbulence: from tracers to intermittency and transport (II)

### Massimo Cencini

Istituto dei Sistemi Complessi CNR Rome, Italy INFN "Tor Vergata", Rome, Italy massimo.cencini@cnr.it







# Outline -topic 2-

- Brief recall of single particle dispersion in turbulence
- Relative dispersion of two particles: Chaos, Richardson dispersion + intermittency, explosive separation
- Some ideas on multiparticle dispersion, shape and size evolution, zero modes

# Single particle: absolute dispersion

$$\dot{\mathbf{X}} = \mathbf{u}(\mathbf{X}, t) + \sqrt{2D_0} \boldsymbol{\eta}(t)$$

We consider small particles with same density as the fluid: no inertia

Gaussian noise

$$\langle \eta_i(t) \rangle = 0 \quad \langle \eta_i(t) \eta_j(t') \rangle = \delta_{ij} \delta(t - t')$$

A simple limiting case

no flow (Brownian diffusion)

$$\begin{split} \dot{\boldsymbol{X}} &= \sqrt{2D_0} \boldsymbol{\eta}(t) \\ X_i(t) - X_i(0) &= \sqrt{2D_0} \int_0^t ds \, \eta_i(s) \\ \langle [X_i(t) - X_i(0)]^2 \rangle &= 2D_0 \int_0^t ds \, \int_0^t ds' \, \langle \eta_i(s) \eta_i(s') \rangle = 2D_0 t \\ &\qquad \qquad \delta(s-s') \end{split}$$

# Single particle: absolute dispersion

Another limit: no noise (only advection)

$$\dot{m{X}} = m{V}(t) = m{u}(m{X}(t),t)$$

$$X_i(t) - X_i(0) = \int_0^t ds \, V_i(s) \qquad \langle [X_i(t) - X_i(0)]^2 \rangle = \int_0^t ds \, \int_0^t ds' \, \langle V_i(s) V_i(s') \rangle$$
by stationarity

Lagrangian Correlation function 
$$\langle V_i(s)V_i(s')\rangle = C_{ii}(s,s') = \widetilde{C_{ii}}(s-s')$$

$$\langle [X_i(t) - X_i(0)]^2 \rangle = \int_0^t ds \int_0^t ds' C_{ii}(s - s') = 2 \int_0^t ds \int_0^s ds' C_{ii}(s - s') = 2 \int_0^t ds \int_0^s d\tau C_{ii}(\tau)$$

 $\tau_{c}$ 

$$\tau_c = \frac{\int_0^\infty d\tau C_{ii}(\tau)}{C_{ii}(0)}$$

Lagrangian correlation time 
$$\tau_{\text{c}} = \frac{\int_0^\infty d\tau C_{ii}(\tau)}{C_{ii}(0)} \qquad ([X_i(t) - X_i(0)]^2) = \begin{cases} C(0)t^2 & t \ll \tau_c \\ 2[C(0)\tau_c]t & t \gg \tau_c \end{cases}$$

$$C(0) = \langle V^2 \rangle \qquad \qquad D^E = C(0)\tau_c$$

**Eddy diffusivity** 

# Single particle: superdiffusion

What does happen if the correlation function is not integrable? i.e. if the correlation time is not finite

$$\tau_c = \frac{\int_0^\infty d\tau C_{ii}(\tau)}{C_{ii}(0)} = \infty$$



**Superdiffusion (Anomalous diffusion)** 

$$\langle [X_i(t) - X_i(0)]^2 \rangle \sim t^{\alpha} \quad \alpha > 1$$

In turbulence the correlation time is finite and thus we do not expect anomalous diffusion for the single particle But the long time diffusive behavior will be much faster than in the absence of flow due to eddy diffusivity  $D^E \sim \tau_c U^2$ 

# Relative dispersion

$$\dot{\boldsymbol{X}}_1 = \boldsymbol{u}(\boldsymbol{X}_1,t)$$
  
 $\dot{\boldsymbol{X}}_2 = \boldsymbol{u}(\boldsymbol{X}_2,t)$ 

$$R = X_2 - X_1$$


$$\dot{\boldsymbol{R}} = \boldsymbol{u}(\boldsymbol{X}_1 + \boldsymbol{R}, t) - \boldsymbol{u}(\boldsymbol{X}_1, t) = \delta_{\boldsymbol{R}} \boldsymbol{u}$$

### Relative dispersion depends on the two-point properties of the velocity field

Different behaviors depending on the initial separation

$$R = |\mathbf{R}|$$

$$R \ll \eta \to \delta_R u \propto R$$
 
$$\Pi \quad \eta \ll R \ll L \to \delta_R u \propto R^{1/3}$$
 
$$R \gg L \to \delta_L u \quad (Gaussian \ uncorrelated \ variables)$$



The third regime is like Brownian diffusion with an enhanced diffusion coefficient what about the other two regimes?

$$R = |\mathbf{R}| \ll \eta$$
  $\dot{\mathbf{R}} = \mathbb{A}\mathbf{R}$   $\mathbb{A}_{ij} = \frac{\partial u_i}{\partial x_j}$   $\nabla \cdot \mathbf{u} = 0 \mapsto Tr[\mathbb{A}] = 0$ 

In 3D even for non turbulent flows (i.e. laminar flows) particles separate exponentially due to chaos!

$$R(t) pprox R(0) e^{\lambda_1 t} \qquad \lambda_1 > 0 \qquad$$
 the maximal Lyapunov Exponent

$$oldsymbol{R}(t) = \mathbb{W}(0,t) oldsymbol{R}(0) \qquad \mathbb{W}(0,t) = \mathcal{T} \exp \left[ \int_0^t \mathbb{A}(s) ds \right]$$

$$\left[ \underbrace{\mathbb{W}^{\dagger}(0,t)\mathbb{W}(0,t)}^{1/2} \right]^{1/2} = \mathbb{V}(\boldsymbol{x}_0,t) = \mathbb{Q}(\boldsymbol{x}_0,t)\mathbb{D}(\boldsymbol{x}_0,t)\mathbb{Q}^{\dagger}(\boldsymbol{x}_0,t) \quad \mathbb{D}(\boldsymbol{x}_0,t) = \operatorname{diag}\{e^{t\gamma_1(\boldsymbol{x}_0,t)},\ldots,e^{t\gamma_d(\boldsymbol{x}_0,t)}\}$$

positive and symmetric

finite time Lyapunov exponents


**Oseledec Theorem** 

$$\gamma_i(\boldsymbol{x}_0,t) \xrightarrow[t \to \infty]{} \lambda_i(\boldsymbol{x}_0) = \lambda_i$$
if ergodic

$$\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_d$$
 Lyapunov exponents

 $\lambda_1 + \lambda_2$  => growth rate of infinitesimal surfaces  $\lambda_1 + \lambda_2 + \lambda_3 =$  growth rate of infinitesimal volumes incompressibility

$$\dot{m{R}}=\mathbb{A}m{R}$$
  $\mathbb{A}_{ij}=rac{\partial u_i}{\partial x_i}$   $m{
abla}\cdotm{u}=0\mapsto Tr[\mathbb{A}]=0$ 





#### conservation of volumes


$$\lambda_1 + \lambda_2 + \lambda_3 = 0$$


Nonlinear effects lead to folding

### **Chaotic stretching**



$$L_1(0) = L_2(0) = \epsilon$$





Stretching and folding are at the base of mixing in laminar flows

### **Finite Time Lyapunov Exponent**

$$\gamma_1(t) = \frac{1}{t} \log \left( \frac{R(t)}{R(0)} \right)$$

#### Law of large numbers (Oseledec)


$$\lim_{t \to \infty} \lim_{R(0) \to 0} \gamma_1(t) = \lambda_1$$

#### Large deviation theory

$$P_t(\gamma) \sim e^{-S(\gamma)t}$$
  
 $S(\gamma) > 0 \quad if \quad \gamma \neq \lambda_1 \quad \& \quad S(\lambda_1) = 0$ 

#### For large times and close to the minimum (TLC)

$$P_t(\gamma) \sim \exp\left[-\frac{t(\gamma - \lambda_1)^2}{2\sigma^2}\right]$$



J. Bec et al. *Physics of Fluids* 18, 091702 (2006).


**K41** D. Ruelle, Phys. Lett. 72A, 81 (1979)

$$\lambda_1 \sim \frac{1}{\tau_\eta} \sim \frac{1}{T_L} \left( \frac{T_L}{\tau_\eta} \right) \sim Re^{1/2} \sim Re_\lambda$$


MF A. Crisanti, M.H. Jensen, A. Vulpiani, & G. Paladin, PRL 70, 166 (1993)

$$\lambda_1 \sim \frac{1}{\tau_{\eta}} \sim \frac{1}{T_L} \left( \frac{T_L}{\tau_{\eta}} \right) \sim Re^{\frac{1-h}{1+h}} \qquad P(h) \sim \left( \frac{\eta}{L} \right)^{3-D(h)} \sim Re^{\frac{D(h)-3}{1+h}}$$

$$\lambda_1 \sim Re^{\chi} \qquad \chi = \max_h \left\{ \frac{D(h)-2-h}{1+h} \right\}$$



A. Crisanti, M.H. Jensen, A. Vulpiani, G. Paladin. PRL 70, 166 (1993)



J. Bec et al. *Physics of Fluids* 18, 091702 (2006).

# Dispersion in the inertial range: Richardson


The history of relative dispersion in turbulence starts with Richardson (1926) predating K41

From experimental observations of the evolution of the separation of balloons in the atmosphere

$$D_{turbo}(R) \sim R^{4/3}$$
 
$$\partial_t P = R^{-2} \partial_R [D(R) R^2 \partial_R P]$$
 
$$R' = \lambda R \qquad t' = \lambda^a t$$
 
$$\lambda^{-a} \partial_{t'} P = \lambda^{-2/3} R'^{-2} \partial_{R'} [D(R') R'^2 \partial_{R'} P]$$
 invariant if  $a = 2/3 \Longrightarrow R \sim t^{3/2}$ 

so we expect

$$\langle R^2(t) \rangle \sim t^3$$



**Fig. 5.** Figure from the original Richardson's paper [8] representing the experimental results of the diffusivity *D* measured as a function of *l*, i.e. distance separation between two particles. Starting from these data, Richardson proposed his famous 4/3 law drawn in the figure. *Source:* Reproduced from [8].

L.F. Richardson, Proc. Roy. Soc. A 110, 709 (1926).

# Dispersion in the inertial range: Richardson

$$\dot{\mathbf{R}} = \mathbf{u}(\mathbf{X}_1 + \mathbf{R}, t) - \mathbf{u}(\mathbf{X}_1, t) = \delta_{\mathbf{R}}\mathbf{u}$$
$$\frac{\partial R^2}{\partial t} = 2\mathbf{R} \cdot \delta_{\mathbf{R}}\mathbf{u} \sim 2R(\epsilon R)^{1/3} = 2\epsilon^{1/3}R^{4/3}$$

**Consistent with K41** 

$$D_{turbo}(R) \sim R^{4/3}$$

$$\partial_t P = R^{-2} \partial_R [D(R)R^2 \partial_R P] \qquad P(R,t) \propto \frac{1}{(\mathcal{D}\epsilon^{1/3}t)^{9/2}} \exp\left(-\frac{9R^{2/3}}{4\mathcal{D}\epsilon^{1/3}t}\right) \qquad \langle R^2(t) \rangle = g\epsilon t^3$$

$$D(R) = \mathcal{D}\epsilon^{1/3}R^{4/3}$$

NB: the underlying assumption of Richardson diffusion equation is time decorrelation

In analogy with single particle dispersion we can write

$$\langle (R(t) - R(0))^2 \rangle = \int_0^t ds \int_0^t ds' \langle \delta_R v(s) \delta_R v(s') \rangle$$

$$\langle R^2(t) \rangle^{2/3} \sim t^2 \sim \frac{\partial \langle R^2 \rangle}{\partial t} = 2 \int_0^t \langle \delta_R v(t) \delta_R v(s) \rangle = 2 \tau_c(t) \langle \delta_R v^2 \rangle$$

$$\langle \delta_R v^2 \rangle \sim \langle R^2(t) \rangle^{1/3} \sim t$$

$$\tau_c(t) \sim \langle R^2(t) \rangle^{1/3} \sim t$$

which implies that  $\delta_{\!R} v$  remains correlated invalidating the approach

G. Falkovich, K. Gawe dzki, M. Vergassola Rev. Mod. Phys. 73, 913 (2001)

### Dispersion in the inertial range: Richardson

### Persistence of correlations


Shnapp, et al (2023). Nature Commu, 14(1), 4195.

$$\dot{\mathbf{R}} = \mathbf{u}(\mathbf{X}_1 + \mathbf{R}, t) - \mathbf{u}(\mathbf{X}_1, t) = \delta_{\mathbf{R}}\mathbf{u}$$

$$\frac{\partial R^2}{\partial t} = 2\mathbf{R} \cdot \boldsymbol{\delta_R} \boldsymbol{u}$$

$$\frac{\partial R}{\partial t} = \frac{\mathbf{R} \cdot \mathbf{\delta_R u}}{R}$$

$$\cos \theta = \frac{\mathbf{R} \cdot \mathbf{\delta_R u}}{R \delta_R u} = \frac{1}{\delta_R u} \frac{\partial R}{\partial t}$$



### When is the diffusive equation valid?

I. M. Sokolov, PRE 60.5 (1999): 5528.

$$\langle (\delta_R v)^2 \rangle \sim (\delta_L v)^2 \left(\frac{R}{L}\right)^{\alpha} \qquad \frac{\alpha}{2} + \beta < 1$$

$$\tau_R \sim T_L \left(\frac{R}{L}\right)^{\beta} \qquad \partial_t P = R^{-2} \partial_R [D(R) R^2 \partial_R P]$$

$$D(R) \sim R^{\alpha + \beta}$$

Richardson 
$$\alpha = \beta = 2/3$$
 is marginal!

Diffusion eq holds if the displacement from R to  $R+\Delta_R$  in a time  $\tau_R$  is much smaller than R

$$\Delta_R \sim \delta_R u \tau_R \sim \delta_L u T_L \left(\frac{R}{L}\right)^{\alpha/2+\beta}$$

So far we ignored the initial separation  $R_0 = R(0)$ 

Typical time at scale R


$$\tau(R) \sim \frac{R}{\delta_R u} \sim T_L \left(\frac{R}{L}\right)^{2/3}$$

So we should expect that if we start with  $R_0 \in [\eta, L]$  the initial velocity  $\delta_{R_0} u$  will be "maintained" for a time  $\tau(R_0)$ 

Therefore  $0 < t < \tau(R_0)$  we should expect a "ballistic" regime  $\langle R^2(t) \rangle \approx \langle (\delta_{R_0} u)^2 \rangle \, t^2$  (so-called Batchelor dispersion)


$$\langle R^2(t)\rangle \approx \langle (\delta_{R_0} u)^2\rangle t^2$$

This adds some difficulties in observing Richardson dispersion for large initial separations



M Bourgoin, NT. Ouellette, H Xu, J. Berg,

E. Bodenschatz Science 311, 835 (2006)



G. Boffetta & I. Sokolov PRL 88, 094501 (2002)

N. Ouellette, et al NJP 8, 109 (2006)

$$\langle |\boldsymbol{R}(t) - \boldsymbol{R}_0|^2 \rangle = \langle (\delta_{R_0} \boldsymbol{u} \cdot \delta_{R_0} \boldsymbol{u}) \rangle t^2 + \langle \delta_{R_0} \boldsymbol{u} \cdot \delta_{R_0} \boldsymbol{a} \rangle t^3 + o(t^3)$$



 $\approx -2\epsilon$  J. Mann, S. Ott, & JS Andersen (1999)

R.J. Hill JoT 7, N43 (2006)

$$t_0 = S_2(r_0)/(2\epsilon)$$

#### time scale for the end of the ballistic regime

The data collapse extends to times larger than  $t_0$  when the mean-squared separation tends to Richardson  $t^3$  regime. This unexpected fact implies that  $t_0$  is not only the time scale of departure from the ballistic regime, but also that of convergence to Richardson's law. More precisely, numerical data suggest that for  $t \gg t_0$ 

$$\langle |\delta \mathbf{x}(t) - \delta \mathbf{x}(0)|^2 \rangle_{r_0} = g \epsilon t^3 \left[ 1 + C t_0 / t \right] + \text{h.o.t.}$$
 (4)

C does not strongly depend on the Reynolds number. Sys-

#### But depends on the initial separation!

$$C = C(R_0)$$

C = 0 for  $R_0 \approx 4\eta$ Optimal choice




FIG. 1. (Color online) Time evolution of the mean-square separation for  $R_{\lambda} = 730$  and various initial separations. The dashed line represents the behavior (2). The solid line is a fit to the Richardson regime (4) with g = 0.52 and C = 1.6. Inset:  $t_0$  as a function of  $r_0$  in dissipative-scale units. The solid line is an Eulerian average, the circles are Lagrangian measurements, and the dashed line is the turnover time  $\tau(r_0)$ .

R. Bitane, H. Homann, J. Bec PRE **86**, 045302(R) (2012)




Figure 2. Time-evolution of the mean-squared distance for  $R_{\lambda} = 730$  (a) and  $R_{\lambda} = 460$  (b) for various initial separations  $r_0$  as labeled. The horizontal and vertical solid lines represent the integral scale L and its associated turnover time  $\tau_L$ , respectively. The dashed line corresponds to the explosive Richardson-Obukhov law (3) with g = 0.52.

R. Bitane, H. Homann, J. Bec JoT **14:2**, 23-45 (2013)

In the EXP only quite large initial separation were available and they could not be followed for a long time that's why only the ballistic (Batchelor) regime was found

### How good is Richardson PDF with respect to data?

$$P(R,t) \propto \frac{R^2}{\langle R^2(t) \rangle^{3/2}} \exp\left(-A \frac{R^{2/3}}{\langle R^2(t) \rangle^{1/3}}\right)$$




FIG. 2. Probability distribution function of relative separations at three different times. The continuous line is the Richardson prediction (3), and the dashed line is the Gaussian distribution proposed by Batchelor.

G. Boffetta & I. Sokolov PRL 88, 094501 (2002)




FIG. 3. Log-lin plot of  $P(r_n, t)$  versus the rescaled variable  $r_n$  (see text) for  $t = (20, 30, 40, 60, 90, 120)\tau_n$ . The distribution  $P(r_n, t)$  has been divided by a factor  $r_n^2$  to highlight the large separation range. The Richardson prediction, Eq. (4), becomes time independent if rescaled in this way (solid curve). Inset: PDFs plotted only for separations  $r_n$  that, at time lag  $t \in [10:120]\tau_n$ , belong to the inertial subrange.

### R. Scatamacchia, L. Biferale & F. Toschi PRL 109, 144501 (2013)





Figure 6. Probability density function of the distance r at time  $t=2.5\,t_0$  (a) and  $t=5\,t_0$  (b) and for various values of the initial separation. We have here normalized it by  $4\pi\,r^2$  and represented on a log y axis as a function of  $r/\langle |{\bf R}(t)|^2\rangle^{1/2}$ . With such a choice, Richardson's diffusive density distribution (2) appears as a straight line (represented here as a black dashed line).

R. Bitane, H. Homann, J. Bec JoT **14:2**, 23-45 (2013)

### A source of difficulties



R. Scatamacchia, L. Biferale & F. Toschi PRL 109, 144501 (2012)

but see also R. Bitane, H. Homann, J. Bec JoT 14:2, 23-45 (2013)

coexistence of pairs separating very fastly and pairs remaining close to each other for a long time this makes very difficult to interpret statistics at fixed times

### Richardson diffusion: are there effects of intermittency?

$$\langle R^p(t) 
angle \sim t^{eta_p} ~~ eta_p = rac{3}{2} p ~~$$
 dimensionally from Richardson

What do we expect according to MF model?

$$\left\langle \frac{dR^p}{dt} \right\rangle = p \langle R^{p-1} \delta_R u \rangle$$

$$\delta_R u \sim \delta_L u \left(\frac{R}{L}\right)^h$$

$$\left\langle \frac{dR^p}{dt} \right\rangle = p \delta_L u L^{p-1} \int dh \left(\frac{R}{L}\right)^{p-1+h+3-D(h)}$$

MF relation between times and scales 
$$t \sim \frac{R}{\delta_R u} \sim T_L \left(\frac{R}{L}\right)^{1-h} \qquad \frac{R}{L} \sim \left(\frac{t}{T_L}\right)^{\frac{1}{1-h}} \qquad \left\langle \frac{dR^p}{dt} \right\rangle = p \frac{L^p}{T_L} \int dh \left(\frac{t}{T_L}\right)^{\frac{p-1+h+3-D(h)}{1-h}}$$

$$\beta_p = 1 + \min_h \left\{ \frac{p - 1 + h + 3 - D(h)}{1 - h} \right\} = \min_h \left\{ \frac{p + 3 - D(h)}{1 - h} \right\}$$

### Richardson diffusion: are there effects of intermittency?

$$\langle R^p(t) 
angle \sim t^{eta_p} \quad eta_p = rac{3}{2} p \quad ext{dimensionally from Richardson}$$
 
$$eta_p = 1 + \min_h \left\{ rac{p-1+h+3-D(h)}{1-h} 
ight\} = \min_h \left\{ rac{p+3-D(h)}{1-h} 
ight\}$$

$$\mathbf{p=2}$$

$$\min_{h} \left\{ \frac{5 - D(h)}{1 - h} \right\} = 3$$

$$\min_{h} \left\{ 3h + 3 - D(h) \right\} = 1$$

$$\beta_2 = 3$$

$$\langle R^2(t) \rangle = g\epsilon t^3$$





FIGURE 9. Multifractal exponents for pair separation statistics, derived from the scaling exponents of the Eulerian longitudinal structure functions,  $D_L(h)$  ( $\bigcirc$ ), and from the scaling exponents of Eulerian transversal structure functions,  $D_T(h)$  ( $\square$ ). The continuous line is the dimensional Richardson scaling,  $\alpha(p) = 3p/2$ .

### Richardson diffusion: are there effects of intermittency?



10<sup>0</sup>  $10^{0}$  $\triangle \mathbf{R}(t) - \mathbf{R}(0)|^{6} \vee L^{6}$  $|\mathbf{R}(t) - \mathbf{R}(0)|^{4} \times |\mathbf{L}^{4}|^{10} \times |\mathbf{L}^{4}|^{10}$  $r_0 = 12\eta$  $10^{-3} 10^{-2} 10^{-1} 10^{0} 10^{1} 10^{0}$  $r_0 = 24\eta$  $_{\triangle}$   $r_0 = 32\eta$ 10  $_{\triangleright}$   $r_0 = 64\eta$ (a) (b)  $r_0 = 96\eta$ 10 10  $10^{-4} < |\mathbf{R}(t) - \mathbf{R}(0)|^2 > / L^2$  $10^{-2}$  $10^{-2}$  $10^{-8}$  $10^{-6}$  $10^{0}$  $10^{-4}$  $10^{0}$  $<|\mathbf{R}(t)-\mathbf{R}(0)|^2>/L^2$ 

Figure 4. (a) Fourth-order moment  $\langle |\mathbf{R}(t) - \mathbf{R}(0)|^4 \rangle$  and (b) sixth-order moment  $\langle |\mathbf{R}(t) - \mathbf{R}(0)|^6 \rangle$  as function of  $t/t_0$  for  $R_{\lambda} = 730$ . Both curves are normalized such that their expected long-time behavior is  $\propto (t/t_0)^6$  and  $\propto (t/t_0)^9$ , respectively. The black dashed lines represent such behaviors.

Figure 5. Fourth (a) and sixth (b) order moments of |R(t) - R(0)| as a function of its second-order moment for  $R_{\lambda} = 730$ . The two gray dashed lines show a scale-invariant behavior, i.e.,  $\langle |R(t) - R(0)|^4 \rangle \propto \langle |R(t) - R(0)|^2 \rangle^2$  and  $\langle |R(t) - R(0)|^6 \rangle \propto \langle |R(t) - R(0)|^2 \rangle^3$ , respectively. The two insets show the associated local slopes, that is the logarithmic derivatives  $d \log \langle |R(t) - R(0)|^p \rangle / d \log \langle |R(t) - R(0$ 

R. Bitane, H. Homann, J. Bec JoT 14:2, 23-45 (2013)

St = 0.0 Richardson 
$$p=6$$
 $10^4$ 
 $10^3$ 
 $10^2$ 
 $10^1$ 
 $10^0$ 
 $10^0$ 
 $10^1$ 
 $10^1$ 
 $10^2$ 
 $10^2$ 

L. Biferale, A.S Lanotte, R. Scatamacchia, & F. Toschi JFM 757 (2014): 550-572.

### Richardson diffusion: intermittency & exit times

A possibility to mitigate the statistical effects of pairs that separates quickly or remains close for a long time is to look at the exit time statistics

Time for R(t) to pass from R to  $\rho R$  with  $\rho > 1$  this eliminates part of the problem with fixed time statistics and focus on the spatial scale avoiding averaging events which at the same time involve very different scales.

$$\mathcal{T}(R) = \{ R(t) = R \quad \& \quad R(t + \mathcal{T}(R)) = \rho R \}$$

Then one can study their statistics

$$\mathcal{T}(R) \sim \frac{R}{\delta_R u} \sim T_L \left(\frac{R}{L}\right)^{1-h}$$

$$\left\langle \frac{1}{\mathcal{T}^p(R)} \right\rangle \sim \frac{1}{T_L^p} \int dh \left(\frac{R}{L}\right)^{-p+\frac{hp+3-D(h)}{2}} \sim \frac{1}{T_L^p} \left(\frac{R}{L}\right)^{\sigma_p}$$

$$\sigma_p = \zeta_p - p$$

K41+Richardson 
$$\sigma_p = -\frac{2}{3}p$$

G. Boffetta & I. Sokolov PRL 88, 094501 (2002)

G. Boffetta, A. Celani, A. Crisanti, A. Vulpiani, PRE 60, 6734 (1999)




FIG. 3. First moments of the inverse doubling time  $\langle [1/T(R)]^p \rangle$  compensated with Kolmogorov scaling  $R^{-2p/3}$ . Deviations from dimensional compensation are evident, in particular for p=4. In the inset we plot the compensated mean doubling time according to (6) together with the estimate corresponding to  $C_2 \approx 0.55$ .




FIG. 4. First moments of the inverse doubling time  $\langle [1/T(R)]^p \rangle$  compensated with best fit exponent  $\beta_p$ . Observe the improvement in the compensation with respect to Fig. 3. In the inset we plot the structure function exponents estimated from  $\zeta_p = p + \beta_p$ . The dashed line represents the Eulerian exponents obtained by ESS analysis.

### Richardson diffusion & role of intermittency

Not completely settled

VOLUME 88, NUMBER 9

PHYSICAL REVIEW LETTERS

4 March 2002

Relative Dispersion in Fully Developed Turbulence: The Richardson's Law and Intermittency Corrections

G. Boffetta1 and I.M. Sokolov2

PRL 109, 144501 (2012)

PHYSICAL REVIEW LETTERS

week ending 5 OCTOBER 2012

Extreme Events in the Dispersions of Two Neighboring Particles Under the Influence of Fluid Turbulence

R. Scatamacchia, 1,2 L. Biferale, 1 and F. Toschi 3

PHYSICAL REVIEW E

VOLUME 60, NUMBER 6

DECEMBER 1999

Pair dispersion in synthetic fully developed turbulence

G. Boffetta and A. Celani

Dipartimento di Fisica Generale, Università di Torino, Via Pietro Giuria 1, 10125 Torino, Italy; Istituto di Cosmogeofisica, c. Fiume 4, 10133 Torino, Italy; and Istituto Nazionale di Fisica della Materia, Unità di Torino, Torino, Italy

A. Crisanti and A. Vulpiani

Dipartimento di Fisica, Università di Roma "La Sapienza," Piazza le Aldo Moro 2, 00185 Roma, Italy and Istituto Nazionale di Fisica della Materia, Unità di Roma I, Rome, Italy

(Received 1 June 1999)

PHYSICAL REVIEW LETTERS 131, 064001 (2023)

Spontaneous Stochasticity in the Presence of Intermittency

André Luís Peixoto Considera 1,2 and Simon Thalabard 3

PHYSICAL REVIEW E 86, 045302(R) (2012)

#### Time scales of turbulent relative dispersion

Rehab Bitane, Holger Homann, and Jérémie Bec

Laboratoire Lagrange, UMR7293, Université de Nice-Sophia Antipolis, CNRS, Observatoire de la Côte d'Azur,

BP 4229, 06304 Nice Cedex 4, France

(Received 29 June 2012; published 31 October 2012)

Journal of Turbulence, 2013

Vol. 14, No. 2, 23–45, http://dx.doi.org/10.1080/14685248.2013.766747



#### Geometry and violent events in turbulent pair dispersion

Rehab Bitane, Holger Homann and Jérémie Bec\*

J. Fluid Mech. (2014), vol. 755, R4, doi:10.1017/jfm.2014.445

### JFM RAPIDS journals.cambridge.org/rapids



Turbulent pair dispersion as a continuous-time random walk

Simon Thalabard<sup>1</sup>, Giorgio Krstulovic<sup>1</sup> and Jérémie Bec<sup>1</sup>,†



### Turbulent pair dispersion as a ballistic cascade phenomenology

Published online by Cambridge University Press: 08 May 2015

Journal of Fluid Mechanics 772 (2015): 678-704.

Show author

### Richardson dispersion & irreversibility


$$\langle |\mathbf{R}(t) - \mathbf{R}_0|^2 \rangle = \langle (\delta_{R_0} \mathbf{u} \cdot \delta_{R_0} \mathbf{u}) \rangle t^2 + \langle \delta_{R_0} \mathbf{u} \cdot \delta_{R_0} \mathbf{a} \rangle t^3 + o(t^3)$$
  
  $\approx -2\epsilon$ 

J. Mann, S. Ott, & JS Andersen (1999) R.J. Hill JoT 7, N43 (2006)

 $\langle |\delta \mathbf{R}(-t)|^2 \rangle - \langle |\delta \mathbf{R}(t)|^2 \rangle = -2 * neg |t|^3 = 4\epsilon t^3$ 

 $\langle |\delta \mathbf{R}(t)|^2 \rangle = pos \ t^2 + neg \ t^3$ 

### Backward separation is faster than forward separation



labeled for DNS calculations at  $Re_{\lambda} = 38$ .

B. L. Sawford et al

PoF 17, 095109 (2005)




FIG. 4. Forwards relative dispersion (solid lines) and backwards relative FIG. 1. (color online). The difference between the backward dispersion (dashed lines) for initial separations  $r_0/\eta = \frac{1}{4}$ , 1, 4, and 16 as and forward mean squared relative separation,  $(\delta \mathbf{R}(-t)^2 \delta \mathbf{R}(t)^2$ , compensated using Eq. (5). The symbols correspond to experiments: circles for  $R_{\lambda} = 690 \ (R_0/\eta = 267, 333, 400)$ . stars for  $R_{\lambda} = 350 \ (R_0/\eta = 152, 182, 212)$ , and squares for  $R_{\lambda} = 270 \ (R_0/\eta = 95, 114, 133)$ . The lines correspond to DNS at  $R_{\lambda} = 300 \ (R_0/\eta = 19, 38, 58, 77, 92, 123).$ 

J Jucha et al. PRL 113.5 (2014): 054501.

105101-8 Buaria, Sawford, and Yeung

Phys. Fluids 27, 105101 (2015)





FIG. 2. Mean-squared relative displacement as a function of forward time (solid lines, in red) and backward time (dashed lines, in blue) at R<sub>A</sub> 140 (left) and 1000 (right), scaled by Kolmogorov variables, for different initial separations. Arrows indicate direction of increasing  $\tilde{r}_0$ , in logarithmically spaced intervals:  $\tilde{r}_0/\eta = 1/4, 1, 4, 16, 64, 256$ , and 1024 for  $R_A$  140;  $\tilde{r}_0/\eta = 1, 4, 16, 64, 256, 1024$ , and 4096 for  $R_A$  1000. (Results at  $\tilde{r}_0/\eta = 1/4$  for  $R_A$  1000 are not shown since they are not well sampled.)

### Richardson dispersion on the field



#### Figure 6

Mean square separation distance versus time obtained from time-shifted float data during the TOPOGULF experiment: (a) western floats and (b) eastern floats.  $r_0 < 7.5$  km for the 10-day (crosses) and 60-day (open diamonds) reinitialized time series. Asterisks give the corresponding evolution for portions selected with the minimum separation criterion. This consists of finding the minimum separation distance over the entire lifetime of the pair and recording the evolution for 120 days thereafter. The remaining record is then searched for a new minimum and the process is repeated. Numbers of pairs are indicated within parentheses.  $\alpha_W$  and  $\alpha_E$  are numerical coefficients, and  $K_E(\infty)$  and  $K_W(\infty)$  are the diffusion coefficients in the diffusive limit. Figure adapted from Ollitrault et al. 2005.

Ollitrault M, Gabillet C, de Verdie re AC. 2005. Open ocean regimes of relative dispersion. J. Fluid Mech. 533:381–407

### Two-Particle Dispersion in Isotropic Turbulent Flows




Fig. 6. Scale-dependent relative diffusivities. Lower left axes: Tracer-based diffusivity estimates based on fitting ellipses. The solid red, green, and blue symbols show Okubo (36) estimates of  $K_a(3\sigma) = \sigma^2/4t$ . Corresponding estimates from the S1 drifter data are shown by solid black squares. Upper right axes: Scale-dependent mixing length diffusivities,  $K_{I}(r) = r\Delta v(r)$ , observed in S1 launch plotted with uncertainty estimates in solid black lines and filled black circles. Richardson–Obukhov scaling law,  $K_L(r) \sim r^{4/3}$  is indicated.

Poje, Andrew C., et al. "Submesoscale dispersion in the vicinity of the Deepwater Horizon spill." PNAS 111.35 (2014): 12693-12698.

Juan P.L.C. Salazar and Lance R. Collins Annu. Rev. Fluid Mech. 2009. 41:405–32

### Richardson diffusion: why so difficult?

### Which are the difficulties in studying Lagrangian dispersion?


1 need to accurately know the spatial statistics of the flow velocity along Lagrangian paths *high-resolution*, *high frequency* 

2 need scale separation to disentangle different dispersion regimes: exponential, ballistic, turbulent,..


3 need to have high statistical accuracy: *long records* along *many Lagrangian* paths

4 need to limit the impact of inhomogeneities (walls, borders), unsteadiness, anisotropies, stratification

#### Finite Re effects are very severe



### **Persistency of correlations**



# Breakdown of the Lagrangian flow

$$\frac{d\boldsymbol{x}}{dt} = \boldsymbol{f}(\boldsymbol{x}) \qquad \boldsymbol{x} \in \mathfrak{R}^d$$

be f a smooth velocity field

if f is continuous with the Lipschitz condition (essentially if f is differentiable)

$$||f(x) - f(y)|| \le K||x - y||$$

The solution exists and is unique

$$p(\boldsymbol{x}, t|\boldsymbol{y}, s) = \delta(\boldsymbol{x} - \boldsymbol{x}(t; \boldsymbol{y}, s))$$

$$\delta \boldsymbol{x}(t) = \boldsymbol{x}'(t) - \boldsymbol{x}(t)$$

$$\delta \boldsymbol{x}(0) = \boldsymbol{x}'(0) - \boldsymbol{x}(0) = \varepsilon$$

in the presence of chaos

$$|\delta \boldsymbol{x}(t)| = \epsilon e^{\lambda_1 t}$$

vet

$$\lim_{\epsilon \to 0} |\delta \boldsymbol{x}(t)| = 0$$

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}) + \sqrt{2D_0}\mathbf{\eta}(t)$$

$$\partial_t p(\boldsymbol{x}, t) + \nabla f(\boldsymbol{x}) p(\boldsymbol{x}, t) - D_0 \Delta p(\boldsymbol{x}, t) = 0$$

$$p(\boldsymbol{x},s) = \delta(\boldsymbol{x} - \boldsymbol{y})$$

$$\lim_{D_0 \to 0} p(\boldsymbol{x}, t | \boldsymbol{y}, s) = \delta(\boldsymbol{x} - \boldsymbol{x}(t; \boldsymbol{y}, s))$$

# Breakdown of the Lagrangian flow

$$rac{dm{x}}{dt}=m{u}(m{x})$$
 be u a non smooth incompressible velocity field e.g.  $\delta_Rm{u}\sim R^{1/3}$ 

#### 1d example

$$\frac{dR}{\partial t} = \delta_R u = R^{1/3}$$

$$\mathcal{P}(\mathbf{x}, t) + \mathbf{u}(\mathbf{x}) \cdot \nabla p(\mathbf{x}, t) - D_0 \Delta p(\mathbf{x}, t) = 0$$


$$\mathcal{P}(\mathbf{x}, \mathbf{x}) \in \delta(\mathbf{x} - \mathbf{y})$$

$$\lim_{R(t)\to \oplus} p\left(\mathbf{z},t|\mathbf{y},s\right) = \mathcal{P}_2(\mathbf{z},t')^2$$
 weak solution of

$$R(t) = \left[\frac{2}{3}(t-s)\right]^{3/2}$$

but for  $\epsilon = 0$  also R(t) = 0 is a solution

the Lagrangian flow breaks a trajectory is no more labeled by its initial condition



# Breakdown of the Lagrangian flow



We shall come back to this explosive separation when discussing the transport of substances in turbulent flows

### A clean framework for (not only) Richardson dispersion

The Kraichnan model (R. H. Kraichnan, PoF 11,11, 945 (1968)).

$$\dot{\boldsymbol{r}} = \boldsymbol{v}(\boldsymbol{r}, t) + \sqrt{2\kappa} \boldsymbol{\eta}$$



v(r,t) zero mean Gaussian velocity field with correlation  $\langle v^i({f r},t)v^j({f r}',t') \rangle = 2\,\delta(t-t')D^{ij}({f r}-{f r}')$ 

$$D^{ij}(\mathbf{r}) = D_0 \delta^{ij} - \frac{1}{2} d^{ij}(\mathbf{r}) \qquad \lim_{\substack{\eta \to 0 \\ L \to \infty}} d^{ij}(\mathbf{r}) = D_1 r^{\xi} \left( (d-1+\xi) \, \delta^{ij} - \xi \frac{r^i r^j}{r^2} \right) \qquad \qquad 0 \leqslant \xi \leqslant 2$$

NB: is an incompressible ensemble of velocities

$$\sum_{i} \partial_{r_i} \langle v^i(\boldsymbol{r}, t) v^j(\boldsymbol{r}', t') \rangle = \sum_{j} \partial_{r'_j} \langle v^i(\boldsymbol{r}, t) v^j(\boldsymbol{r}', t') \rangle = 0 \qquad \sum_{i} \partial_{r_i} D^{ij}(\boldsymbol{r} - \boldsymbol{r}') = \sum_{j} \partial_{r_j} D^{ij}(\boldsymbol{r} - \boldsymbol{r}') = 0$$

the delta correlation makes the dynamics of particles reversible

B. Shraiman & E. D. Siggia. Nature 405, 639 (2000) G. Falkovich, K. Gawędzki, M. Vergassola. RMP 73, 913 (2001)

### A clean framework for (not only) Richardson dispersion

$$\langle v^{i}(\mathbf{r},t)v^{j}(\mathbf{r}',t')\rangle = 2\,\delta(t-t')D^{ij}(\mathbf{r}-\mathbf{r}')$$

$$D^{ij}(\mathbf{r}) = D_{0}\delta^{ij} - \frac{1}{2}d^{ij}(\mathbf{r}) \quad \lim_{\substack{\eta \to 0 \\ L \to \infty}} d^{ij}(\mathbf{r}) = D_{1}r^{\xi} \left( (d-1+\xi)\,\delta^{ij} - \xi\,\frac{r^{i}r^{j}}{r^{2}} \right)$$

Two particles joint probability at two different times

obeys the equation 
$$\langle p(\mathbf{r}_1,s;\mathbf{R}_1,t|\mathbf{v})p(\mathbf{r}_2,s;\mathbf{R}_2,t|\mathbf{v})\rangle = \mathcal{P}_2(\mathbf{r}_1,\mathbf{r}_2;\mathbf{R}_1,\mathbf{R}_2;t-s) \\ \mathcal{M}_2 = \sum_{n,n'=1}^2 D^{ij}(\mathbf{r}_n-\mathbf{r}_{n'})\nabla_{r_n^i}\nabla_{r_{n'}^j} + \kappa \sum_{n=1}^2 \nabla_{\mathbf{r}_n}^2 \nabla_{\mathbf{r}_n^i}^2 \nabla$$

relative motion  $r = r_1 - r_2$  and  $R = R_1 - R_2$  $\xi = 2$   $\lim \mathcal{P}(r;R;t) = \delta(R)$  $0 \le \xi < 2 \qquad \lim_{r \to 0} \mathcal{P}(r; R; t) \propto \frac{R^{d-1}}{|t|^{d/(2-\xi)}} \exp\left[-\operatorname{const} \times \frac{R^{2-\xi}}{|t|}\right]$  $(\partial_t - M)\mathcal{P}(r; R, t) = \delta(t)\delta(R - r)$  $M = \frac{1}{r^{d-1}} \partial_r [(d-1)D_1 r^{d-1+\xi} + 2\kappa r^{d-1}] \partial_r$ 

$$\langle R^2(t) \rangle \propto t^{2/(2-\xi)}$$

B. Shraiman & E. D. Siggia. *Nature* 405, 639 (2000) G. Falkovich, K. Gawedzki, M. Vergassola. RMP 73, 913 (2001)

# Multi-particle separation





# Multi-particle separation

N particles 
$$\dot{\boldsymbol{r}}_i = \boldsymbol{v}(\boldsymbol{r}_i,t) + \sqrt{2\kappa}\boldsymbol{\eta}_i \qquad i=1,\ldots,N$$

### **Multiparticle propagator**

$$\left\langle \prod_{n=1}^{N} p(\mathbf{r}_{n}, s; \mathbf{R}_{n}, t | \mathbf{v}) \right\rangle_{\mathbf{v}} = \mathcal{P}_{N}(\underline{\mathbf{r}}; \underline{\mathbf{R}}; t - s)$$

The relative motion of N particles may be described by the versions of the joint PDF's (65) averaged over rigid translations:

$$\widetilde{\mathcal{P}}_{N}(\underline{\mathbf{r}};\underline{\mathbf{R}};t) = \int \mathcal{P}_{N}(\underline{\mathbf{r}};\underline{\mathbf{R}}+\boldsymbol{\varrho};t)d\boldsymbol{\rho}, \tag{66}$$


where  $\boldsymbol{\rho} = (\boldsymbol{\rho}, \dots, \boldsymbol{\rho})$ . The PDF in Eq. (66) describes the distribution of the particle separations  $\mathbf{R}_{nm} = \mathbf{R}_n - \mathbf{R}_m$  or of the relative positions  $\mathbf{\tilde{R}} = (\mathbf{R}_1 - \mathbf{\bar{R}}, \dots, \mathbf{R}_N - \mathbf{\bar{R}})$ .

N=2 we saw that the separation R grows as a power law

$$R \sim t^{3/2}$$
 turbulence  $R \sim t^{1/(2-\xi)}$  Kraichnan

When N>2 besides the evolution of the size we also have evolution of the shape to account for

# Multi-particle separation: e.g. N=4



(Chertkov, Pumir, Shraiman 1999)

$$\rho_0 = (\mathbf{x_1} + \mathbf{x_2} + \mathbf{x_3} + \mathbf{x_4})/2 
\rho_1 = (\mathbf{x_1} - \mathbf{x_2})/\sqrt{2} 
\rho_2 = (\mathbf{2x_3} - \mathbf{x_2} - \mathbf{x_1})/\sqrt{6} 
\rho_3 = (\mathbf{3x_4} - \mathbf{x_3} - \mathbf{x_2} - \mathbf{x_1})/\sqrt{12}$$

It can be studied in terms of the eingenvalues "g" of the inertia matrix  $I = \rho \rho^T$ built from vectors  $\rho_1$ ,  $\rho_2$ ,  $\rho_3$  in the set of coordinates independent of centre of mass  $\rho_0$ 


• 
$$r=\sqrt{g_1+g_2+g_3}=\sqrt{rac{1}{8}\sum_{ij}|m{x}_i-m{x}_j|}$$
 tetrad size

$$ullet I_i \equiv rac{g_i}{r^2}, \;\; I_1 + I_2 + I_3 \equiv 1 \;\;\;$$
 tetrad shape


$$I_1=I_2=I_3=1/3 \ regular$$
 $I_3=0 \ coplanar$ 
 $I_2=I_3=0 \ collinear$ 

$$I_3 = 0$$
 coplanar

$$I_2 = I_3 = 0$$
 collinear







# Multi-particle separation: N=4




FIG. 1. Evolution of the mean eigenvalues  $g_1$  (+),  $g_2$  (×) and  $g_3$  (\*) of the moment of inertia matrix  $I = \rho \rho^T$ . The line represents the dimensional scaling  $t^3$ . In the inset, from top to bottom: evolution at small times of  $\langle \ln A(t) \rangle$  (surface),  $\langle \ln R(t) \rangle$  (distance),  $\langle \ln V(t) \rangle$  (volume). The linear slopes of the three curves in the range of times  $\tau_{\eta} < t < 3\tau_{\eta}$  yield  $\lambda_1 + \lambda_2$ ,  $\lambda_1$ , and  $\lambda_1 + \lambda_2 + \lambda_3$ , respectively.





FIG. 4: Probability density function of shape indices  $I_2$  and  $I_3$  (inset) at times  $t=35\tau_\eta$  (+) and  $t=63\tau_\eta$  (×). The full lines are the pdfs for independent, Gaussian distributed particle positions.

$$\langle I_1 \rangle = 0.854 \quad \langle I_2 \rangle = 0.135 \quad \langle I_3 \rangle = 0.011$$

In general it was found  $I_2 \ll I_1$  and  $I_3 \ll I_2$  denoting preference for elongated and planar geometries


L. Biferale; G. Boffetta; A. Celani; B. J. Devenish; A. Lanotte; F. Toschi PoF 17, 111701 (2005) see also A. Pumir, B.I. Shraiman, M. Chertkov PRL 85, 5324 (2000) for a previous study at lower Re

### Time evolution of tetrads in 3D turbulence




**Courtesy of Guido Boffetta** 

L. Biferale; G. Boffetta; A. Celani; B. J. Devenish; A. Lanotte; F. Toschi PoF 17, 111701 (2005)




# Multi-particle separation: N=4



### **EXP** are in qualitative agreement with DNS

#### **But:**

- 1. no evidence of  $t^3$  size growth
- 2. difficulty to have long time information due to finite volume effects in the experiment
- 3. size of initial tetraedon is quite large



**Figure 5.** The evolution of the PDF of the shape factor  $\langle I_3 \rangle$  for tetrads with an initial size of  $R_0 = 20$  mm. The legends show the time (in units of  $\tau_n$ ) at which the PDFs are measured. Similar changes are observed for tetrads with initial size of 10 and 15 mm. (a)  $R_{\lambda} = 690 (\eta = 30 \,\mu\text{m})$ ; (b)  $R_{\lambda} = 815 (\eta = 23 \,\mu\text{m})$ .

### Multi-particle separation: Lagrangian average

Let us reconsider the Kraichnan model

$$\dot{\boldsymbol{r}} = \boldsymbol{v}(\boldsymbol{r},t) + \sqrt{2\kappa}\boldsymbol{\eta} \quad \langle v^i(\boldsymbol{r},t)v^j(\boldsymbol{r}',t')\rangle = 2\,\delta(t-t')D^{ij}(\boldsymbol{r}-\boldsymbol{r}')$$

$$D^{ij}(\mathbf{r}) = D_0 \delta^{ij} - \frac{1}{2} d^{ij}(\mathbf{r}) \qquad \lim_{\substack{\eta \to 0 \\ I = r \\ N}} d^{ij}(\mathbf{r}) = D_1 r^{\xi} \left( (d - 1 + \xi) \delta^{ij} - \xi \frac{r^i r^j}{r^2} \right) \qquad \qquad 0 \le \xi \le 2$$

smooth

For  $\xi = 0$  (i.e. rough flow) particles undergo independent Brownian motions in the limit  $\kappa \to 0$ 

$$\mathbf{R}_{ij} = \mathbf{x}_i - \mathbf{x}_j \quad \langle R_{ij}^2(t) \rangle = r_{ij}^2 + D_1 t \quad r_{ij} = R_{ij}(0)$$

Now consider N=4 particles and functions,  $f(m{R}_{12},m{R}_{3,4})$  , of their relative separation

e.g. 
$$f(\boldsymbol{R}_{12},\boldsymbol{R}_{3,4}) = R_{12}^2 - R_{3,4}^2$$

Clearly if we average along the Lagrangian trajectories  $\langle f \rangle(t) = \langle f(\boldsymbol{R}_{12}(t), \boldsymbol{R}_{34}(t)) \rangle = \langle R_{12}^2(t) - R_{34}^2(t) \rangle = r_{12}^2 - r_{34}^2 = const$ 

$$f(\mathbf{P}_{-}, \mathbf{P}_{-}) = 2(d+2)P^{2}P^{2}$$
  $d(P^{4}+P^{4})$ 

 $f(\mathbf{R}_{12}, \mathbf{R}_{34}) = 2(d+2)R_{12}^2R_{34}^2 - d(R_{12}^4 + R_{34}^4)$ the same happens to

An example for smooth flows  $\xi=2$   $f({m R}_{12},{m R}_{34})=R_{12}^2/R_{34}^2$  indeed  $R_{ij}=r_{ij}e^{\lambda t}$ 

When this happens such functions are called zero modes

How does it work?

# Multi-particle separation & zero modes

Be  $m{R}=(m{x}_1,m{x}_2,\ldots,m{x}_N)$  and  $m{ ilde{R}}$  the restriction to variables only depending on relative separations of N particles

Lagrangian average

$$\langle f \rangle (\underline{\tilde{r}}, t) = \langle f(\underline{\tilde{R}}(t; \underline{\tilde{r}})) \rangle = \int f(\underline{\tilde{R}}') \, \tilde{\mathcal{P}}(\underline{\tilde{r}}; \underline{\tilde{R}}'; t) d\underline{\tilde{R}}'$$

$$\frac{d\langle f\rangle(\underline{\tilde{\boldsymbol{r}}},t)}{dt} = \int f(\underline{\tilde{\boldsymbol{R}}}') \,\partial_t \tilde{\mathcal{P}}_N(\underline{\tilde{\boldsymbol{r}}};\underline{\tilde{\boldsymbol{R}}}';t)) d\underline{\tilde{\boldsymbol{R}}}' = \int f(\underline{\tilde{\boldsymbol{R}}}') \, \tilde{\mathcal{M}}_N \tilde{\mathcal{P}}_N(\underline{\tilde{\boldsymbol{r}}};\underline{\tilde{\boldsymbol{R}}}';t)) d\underline{\tilde{\boldsymbol{R}}}'$$

Kraichnan Model

$$\partial_t \tilde{\mathcal{P}}_N = \tilde{\mathcal{M}}_N \tilde{\mathcal{P}}_N$$

$$\widetilde{\mathcal{M}}_{N} = -\sum_{n \leq m} \left[ d^{ij}(\mathbf{r}_{nm}) + 2 \kappa \delta^{ij} \right] \nabla_{r_{n}^{i}} \nabla_{r_{m}^{j}}$$

is self-adjoint

$$\frac{d\langle f\rangle(\tilde{\underline{r}},t)}{dt} = \int \tilde{\mathcal{M}}_N f(\tilde{\underline{R}}') \, \tilde{\mathcal{P}}_N(\tilde{\underline{r}};\tilde{\underline{R}}';t)) d\tilde{\underline{R}}'$$

$$\tilde{\mathcal{M}}_N Z_N(\underline{\tilde{\boldsymbol{R}}}) = 0 \Longrightarrow \frac{d\langle Z_N \rangle (\underline{\tilde{\boldsymbol{r}}}, t)}{dt} = 0$$

$$\xi = 0 \quad \tilde{\mathcal{M}}_N \to \quad \begin{array}{c} \text{multidimensional} \\ \text{Laplacian} \end{array} \qquad Z_N \to \quad \begin{array}{c} \text{Harmonic} \\ \text{Polynomials} \end{array}$$

e.g. 
$$R_{12}^2-R_{3,4}^2 \\ N=4 \quad 2(d+2)R_{12}^2R_{34}^2-d(R_{12}^4+R_{34}^4)$$

# Zero modes are non-typical functions

Be  $m{R}=(m{x}_1,m{x}_2,\dots,m{x}_N)$  and  $m{ ilde{R}}$  the restriction to variables only depending on relative separations of N particles

N particles are characterized by the size  $\ R$  and the shape  $\dfrac{{m R}}{R}$ 

#### **Kraichnan Model**

$$\partial_{t} \tilde{\mathcal{P}}_{N} = \tilde{\mathcal{M}}_{N} \tilde{\mathcal{P}}_{N}$$

$$\tilde{\mathcal{M}}_{N} = -\sum_{n < m} \left[ d^{ij}(\mathbf{r}_{nm}) + 2\kappa \delta^{ij} \right] \nabla_{r_{n}^{i}} \nabla_{r_{m}^{j}}$$

$$\sim r_{nm}^{\xi}$$

$$[\tilde{\mathcal{M}}_{N}] = [L]^{\xi - 2} \Longrightarrow [T] \sim [L]^{2 - \xi}$$

$$\mathcal{P}_N(\underline{\tilde{R}},t) = \lim_{r \to 0} \mathcal{P}_N(\underline{\tilde{r}};\underline{\tilde{R}};t) \qquad 0 \le \xi < 2$$

$$\mathcal{P}_N(\underline{\tilde{\mathbf{R}}},t) = \lambda^{(N-1)d} \, \mathcal{P}_N(\lambda \underline{\tilde{\mathbf{R}}}, \lambda^{2-\xi}t)$$

Be f a scaling function

$$f(\lambda \underline{\tilde{R}}) = \lambda^{\sigma} f(\underline{\tilde{R}})$$

$$\begin{split} \langle f \rangle &= \int f(\underline{\tilde{\boldsymbol{R}}}) \mathcal{P}_N(\underline{\tilde{\boldsymbol{R}}},t) d\underline{\tilde{\boldsymbol{R}}} = \lambda^{-\sigma} \int f(\lambda\underline{\tilde{\boldsymbol{R}}}) \lambda^{(N-1)d} \mathcal{P}_N(\lambda\underline{\tilde{\boldsymbol{R}}},\lambda^{2-\xi}t) d\underline{\tilde{\boldsymbol{R}}} \\ &= \lambda^{-\sigma} \int f(\lambda\underline{\tilde{\boldsymbol{R}}}) \mathcal{P}_N(\lambda\underline{\tilde{\boldsymbol{R}}},\lambda^{2-\xi}t) d\underline{\tilde{\boldsymbol{\lambda}}}\underline{\boldsymbol{R}} = \lambda^{-\sigma} \int f(\underline{\tilde{\boldsymbol{R}}}) \mathcal{P}_N(\underline{\tilde{\boldsymbol{R}}},\lambda^{2-\xi}t) d\underline{\tilde{\boldsymbol{R}}} \\ &= t^{\frac{\sigma}{2-\xi}} \int f(\underline{\tilde{\boldsymbol{R}}}) \mathcal{P}_N(\underline{\tilde{\boldsymbol{R}}},1) d\underline{\tilde{\boldsymbol{R}}} \end{split}$$

depends only on the shape

A generic scaling function grows in time a là Richardson in order to be a zero mode the shape evolution should compensante for the size growth

# Multi-particle separation

In the Kraichnan model zero modes exist also for  $\xi \neq 0/2$ 

And are expected to exist also for generic velocity fields (i.e. non reversible, non Markovian and non Gaussian, though for such (general velocities) nobody knows the operator

In non-smooth flows the combination of explosive separation and zero modes has important consequences on the transport of substances

# Summary and take home messages

- Single particle dispersion is typically diffusive unless long anomalous correlations are present
- two (N=2) or many particle (N>2) dispersion probe the spatial statistics of velocity field
- N=2 particle is qualitatively fitting Richardson picture though several difficulties make its experimental and numerical observation non trivial especially when interested to high order statistics
- While in smooth velocity fields the Lagrangian flow is preserved in rough velocity fields one has explosive separation, the Lagrangian flow breaks and become spontaneously stochastic
- N>2 particle dispersion leads to account not only for the evolution of the size but also of the shape which is in general non trivial
- There is a clean framework (The Kraichnan model) where both N=2 and N>2 particle dispersion can be studied
- The study of the Kraichnan model highlights the existence of zero modes: function of relative separation of the particles statistically conserved by the Lagrangian dynamics

# Bibliography

#### **Reviews**

Sawford, Brian. "Turbulent relative dispersion." *Annual review of fluid mechanics* 33.1 (2001): 289-317.

Salazar, Juan PLC, and Lance R. Collins. "Two-particle dispersion in isotropic turbulent flows." *Annual review of fluid mechanics* 41.1 (2009): 405-432. Falkovich, Gregory, K. Gawędzki, and Massimo Vergassola. "Particles and fields in fluid turbulence." *Reviews of modern Physics* 73.4 (2001): 913.

#### Theory/Modeling

Richardson, Lewis Fry. "Atmospheric diffusion shown on a distance-neighbour graph." *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 110.756 (1926): 709-737.

Batchelor, George Keith. "Diffusion in a field of homogeneous turbulence: II. The relative motion of particles." *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 48. No. 2. Cambridge University Press, 1952.

Kraichnan, Robert H. "Dispersion of particle pairs in homogeneous turbulence." *The Physics of Fluids* 9.10 (1966): 1937-1943.

Boffetta, Guido, et al. "Pair dispersion in synthetic fully developed turbulence." *Physical Review E* 60.6 (1999): 6734.

Sokolov, I. M. "Two-particle dispersion by correlated random velocity fields." *Physical Review E* 60.5 (1999): 5528.

Rast, Mark Peter, and Jean-François Pinton. "Pair dispersion in turbulence: the subdominant role of scaling." *Physical review letters* 107.21 (2011): 214501.

Thalabard, S, G Krstulovic, and J Bec. "Turbulent pair dispersion as a continuous-time random walk." *Journal of Fluid Mechanics* 755 (2014): R4.

Eyink, Gregory L., and Dmytro Bandak. "Renormalization group approach to spontaneous stochasticity." *Physical Review Research* 2, no. 4 (2020): 043161.

A. L Peixoto Considera, and S Thalabard. "Spontaneous stochasticity in the presence of intermittency." *Physical Review Letters* 131.6 (2023): 064001.

### **Experiments**

Jullien, Marie-Caroline, Jérôme Paret, and Patrick Tabeling. "Richardson pair dispersion in two-dimensional turbulence." *Physical review letters* 82.14 (1999): 2872.

Ott, Søren, and Jakob Mann. "An experimental investigation of the relative diffusion of particle pairs in three-dimensional turbulent flow." *Journal of Fluid Mechanics* 422 (2000): 207-223.

Bourgoin, Mickaël, et al. "The role of pair dispersion in turbulent flow." Science 311.5762 (2006): 835-838.

Ouellette, Nicholas T., et al. "An experimental study of turbulent relative dispersion models." New Journal of Physics 8.6 (2006): 109.

Berg, Jacob, et al. "Backwards and forwards relative dispersion in turbulent flow: an experimental investigation." *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics* 74.1 (2006): 016304.

Shnapp, Ron, Stefano Brizzolara, Marius M. Neamtu-Halic, Alessandro Gambino, and Markus Holzner. "Universal alignment in turbulent pair dispersion." *Nature Communications* 14, no. 1 (2023): 4195.

Jucha, Jennifer, et al. "Time-reversal-symmetry breaking in turbulence." *Physical review letters* 113.5 (2014): 054501.

# **Bibliography**

#### **DNS-Richardson**

Boffetta, Guido, and Igor M. Sokolov. "Relative dispersion in fully developed turbulence: the Richardson's law and intermittency corrections." *Physical review letters* 88.9 (2002): 094501.

Ishihara, T., & Kaneda, Y. (2002). Relative diffusion of a pair of fluid particles in the inertial subrange of turbulence. *Physics of Fluids*, *14*(11), L69-L72.

Boffetta, Guido, and Igor M. Sokolov. "Statistics of two-particle dispersion in two-dimensional turbulence." *Physics of fluids* 14.9 (2002): 3224-3232.

Yeung, P. K. & Borgas, M. S. (2004). Relative dispersion in isotropic turbulence. Part 1. Direct numerical simulations and Reynolds-number dependence. *Journal* 

Yeung, P. K., & Borgas, M. S. (2004). Relative dispersion in isotropic turbulence. Part 1. Direct numerical simulations and Reynolds-number dependence. *Journal of Fluid Mechanics*, 503, 93-124.

Biferale, L., Boffetta, G., Celani, A., Devenish, B. J., Lanotte, A., & Toschi, F. (2005). Lagrangian statistics of particle pairs in homogeneous isotropic turbulence. *Physics of Fluids*, *17*(11).

Bitane, R., Homann, H., & Bec, J. (2012). Time scales of turbulent relative dispersion. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 86(4), 045302.

Bitane, R., Homann, H., & Bec, J. (2013). Geometry and violent events in turbulent pair dispersion. *Journal of Turbulence*, 14(2), 23-45.

Biferale, L., Lanotte, A. S., Scatamacchia, R., & Toschi, F. (2014). Intermittency in the relative separations of tracers and of heavy particles in turbulent flows. *Journal of fluid mechanics*, 757, 550-572.

Buaria, D., Sawford, B. L., & Yeung, P. K. (2015). Characteristics of backward and forward two-particle relative dispersion in turbulence at different Reynolds numbers. *Physics of Fluids*, 27(10).

#### **Multi-Particle**

Bernard, Denis, Krzysztof Gawedzki, and Antti Kupiainen. "Slow modes in passive advection." *Journal of Statistical Physics* 90, no. 3 (1998): 519-569. Biferale, L., Boffetta, G., Celani, A., Devenish, B. J., Lanotte, A., & Toschi, F. (2005). Multiparticle dispersion in fully developed turbulence. *Physics of Fluids*, *17*(11).

Xu, H., Ouellette, N. T., & Bodenschatz, E. (2008). Evolution of geometric structures in intense turbulence. *New Journal of Physics*, *10*(1), 013012. Pumir, Alain, Eberhard Bodenschatz, and Haitao Xu. "Tetrahedron deformation and alignment of perceived vorticity and strain in a turbulent flow." *Physics of Fluids* 25, no. 3 (2013).

Lüthi, B., Ott, S., Berg, J., & Mann, J. (2007). Lagrangian multi-particle statistics. *Journal of Turbulence*, (8), N45.

Polanco, J. I., Arun, S., & Naso, A. (2023). Multiparticle Lagrangian statistics in homogeneous rotating turbulence. *Physical Review Fluids*, *8*(3), 034602. Hackl, J. F., Pui-Kuen Yeung, and B. L. Sawford. "Multi-particle and tetrad statistics in numerical simulations of turbulent relative dispersion." *Physics of Fluids* 23, no. 6 (2011).

Mazzitelli, I. M., Toschi, F., & Lanotte, A. S. (2014). An accurate and efficient Lagrangian sub-grid model. *Physics of Fluids*, 26(9).

Mazzitelli, I. M., F. Fornarelli, A. S. Lanotte, and P. Oresta. "Pair and multi-particle dispersion in numerical simulations of convective boundary layer turbulence." *Physics of Fluids* 26, no. 5 (2014).