
Rotating turbulence A nice review: Godeferd & Moisy (2015)




$$\frac{\partial t}{\partial u} + \Omega \cdot \nabla u = -2\Omega \times u - \Delta b + 0 \Delta_{\alpha} u$$

$$\frac{1}{2} \frac{1}{2} \frac{1}$$

$$\Rightarrow \partial_{t} \hat{\mathcal{Q}}_{k} = -\sum_{p+q=k} (\hat{\mathcal{Q}}_{p}.iq) \hat{\mathcal{Q}}_{q} - 2\Omega \times \hat{\mathcal{Q}}_{k} - ik\hat{p}_{k} - \nu k^{2}\hat{\mathcal{Q}}_{k}$$

$$p+q=k \qquad (1)$$



$$\hat{e}_{1} = \hat{e}_{1} = \hat{e}_{2} = \frac{k}{|k|}$$

$$\hat{e}_{2} = \frac{k \times \Omega}{|k \times \Omega|}$$

$$\hat{e}_{3} = \hat{e}_{1} \times \hat{e}_{2}$$

$$\hat{e}_{2}$$

$$\hat{e}_{3}$$

$$\hat{e}_{3}$$

$$i k \times (\hat{e}_3 + i s \hat{e}_2) = i k (\hat{e}_4 \times \hat{e}_3 + i s \hat{e}_1 \times \hat{e}_2) =$$

$$= i k (-\hat{e}_2 + i s \hat{e}_3) = -k (s \hat{e}_3 + i \hat{e}_2)$$

$$= -s k (\hat{e}_3 + i s \hat{e}_2)$$

$$U = \sum_{k} \sum_{s=\pm 1}^{\infty} b_{s}(k) h_{s}(k) e^{i(k \cdot x - \sigma_{s}t)}$$

$$R_0 = \frac{U}{2L\Omega} = \frac{Z\Omega}{G} \ll 1 \Rightarrow \frac{Z\Omega}{G} \ll Z_0$$

$$\frac{1}{G} = \frac{4}{U}$$

$$S = \sum_{k, s_k} b_{s_k}(k, t) h_{s_k}(k) e^{i(k \cdot x - \sigma_s t)} = \sum_{k, s_k} b_{s_k}(k, t) h_{s_k}(k) e^{ik \cdot x}$$

$$= \sum_{k, s_k} a_{s_k}(k, t) h_{s_k}(k) e^{ik \cdot x}$$

$$h_{sk}(\underline{e}) \partial_{t} \partial_{s_{k}}(\underline{e}) = - \underbrace{\sum_{k=p+q} (h_{s_{k}}(\underline{e}) \cdot iq) h_{s_{q}}(q) \partial_{s_{p}}(\underline{p}) \partial_{s_{q}}(\underline{q})}_{h_{s_{k}}(\underline{e}) \partial_{s_{k}}(\underline{e}) - i\underline{k} \hat{p}_{\underline{k}}}$$

$$-2\underline{\Sigma} \times h_{s_{k}}(\underline{e}) \partial_{s_{k}}(\underline{e}) - \underline{V}\underline{k}^{2} h_{s_{k}}(\underline{k}) \partial_{s_{k}}(\underline{e}) - i\underline{k} \hat{p}_{\underline{k}}$$

$$\frac{\partial}{\partial t} + Vk^{2}b_{sk}(k) = \frac{1}{2} \sum_{k+p+q=0}^{\infty} C_{kpq} b_{sp}^{*}(P)b_{sq}^{*}(q) e^{i(\sigma_{sk} + \sigma_{s} + \sigma_{s})t}$$

$$\sum_{k+p+q=0}^{\infty} S_{p,s}^{*}(P)b_{sq}^{*}(q) e^{i(\sigma_{sk} + \sigma_{s} + \sigma_{s})t}$$

$$\left(\frac{\partial}{\partial t} + V k^2\right) b_{sk}(k) = \frac{1}{2} \sum_{\substack{k \neq p \neq q = 0}} C_{kpq} b_{sp}^*(\frac{p}{p}) b_{sq}^*(\frac{q}{q})$$


$$\int O_{s_R} = 2s \Omega \cdot \hat{k} = 2s \Omega \cos \Theta_k$$


$$| k \cdot \Omega = k \Omega \cos \Theta_k$$

$$|z \cos \Theta_k + p \cos \Theta_p + q \cos \Theta_q = 0$$

$$|z \cos \Theta_k + z \cos \Theta_p + z \cos \Theta_q = 0$$

$$\frac{\cos \Theta_{k}}{S_{p}q-S_{q}p} = \frac{\cos \Theta_{p}}{S_{q}k-S_{k}q} = \frac{\cos \Theta_{q}}{S_{k}p-S_{p}k}$$









FIG. 2. Isocontours of the axisymmetric energy spectrum $e(k_{\perp}, k_{\parallel})/\sin(\theta_k)$ in the runs with $\Omega = 4$ (above) and 8 (below); dark means larger energy density (in logarithmic scale). Lines indicating the modes for which the wave time becomes equal to the sweeping time, and to the turnover time, are given as references. It should be noted that the energy does not accumulate near the modes with $\tau_{\omega} = \tau_{\rm NL}$, unlike what is expected in theories dealing with the concept of critical balance.²⁴

Phenomenology:




Phenomenology:


$$S_{\frac{1}{2}} = S_{\frac{1}{2}} + S_{\frac{1}{2}}$$


$$S_{\frac{1}{2}} = S_{\frac{1}{2}} + S_{\frac{1}{2}} +$$


$$E(k_1) \sim \frac{S_{0e}}{k_1} \sim \Omega'/2 e^{1/2} k_1^{-2}$$

$$\Rightarrow E(k_1) = C \Omega'/2 e^{1/2} k_1^{-2}$$













FIGURE 5. Angular distribution of energy spectra for different co-latitudes θ , averaged for $5 \le t \le 6$; the Zeman wavenumber is indicated with an arrow. Slopes -7/2 (corresponding to $\sim k^{-5/2}$ scaling in units of the reduced energy spectra) and -8/3 (corresponding to Kolmogorov $\sim k^{-5/3}$ scaling) are shown as a reference. Note the recovery of isotropy beyond k_{Ω} .

$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p - N\theta \hat{z} + \nu \nabla^2 \mathbf{u} + \mathbf{F},$


STABLY STRATIFIED TURBULENCE

$$\partial_t \theta + \mathbf{u} \cdot \nabla \theta = N u_z + \kappa \nabla^2 \theta,$$





2048³ Rorai et al., PRE (2014)

STABLY STRATIFIED TURBULENCE



Stratified turbulence





Slow modes

$$\frac{U_1}{L_1} \approx \frac{U_{11}}{L_1} \implies U_{11} \approx \frac{L_{11}}{L_1} U_1 \ll U_1$$

$$\begin{vmatrix}
\frac{\partial U}{\partial t} + U \cdot \nabla U = -\frac{1}{P} \nabla P - N\Theta^{\frac{2}{2}} + V \nabla^{2} U \\
\frac{\partial O}{\partial t} + U \cdot \nabla O = NW + K \nabla^{2}O
\end{vmatrix}$$
(1)

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - N \nabla \Theta \times \hat{z} + V \nabla^2 \omega$$

$$\frac{\partial \omega}{\partial t} + \omega \cdot \nabla \omega = \omega \cdot \nabla \omega - V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega + V \nabla \omega = \omega + V \nabla \omega +$$

$$\Rightarrow \frac{U_1^2}{L_1 L_1} \sim \frac{N^2 U_{11}}{U_1}$$


$$\Rightarrow Fr_{11}^2 = \frac{U_1^2}{N^2 L_{11}^2} \sim \frac{1}{N^2 L_{11}^2} \frac{N^2 L_1}{V_1} \frac{U_{11}}{U_1} \sim \frac{L_1}{L_{11}} \frac{U_{11}}{U_1} \sim \Lambda$$


$$\Rightarrow Fr_{11} \sim \Lambda$$


$$\Rightarrow Fr_{11} \sim \Lambda$$


$$\Rightarrow \frac{U_1^2}{N^2 L_{11}^2} \sim \frac{1}{N^2 L_{11}^2} \frac{U_1}{V_1} \sim \frac{L_1}{U_1} \frac{U_{11}}{U_1} \sim \Lambda$$

$$Fr_{\perp} = \frac{U_{\perp}}{NL_{\perp}} \sim \frac{L_{\parallel}}{L_{\perp}}$$









Partides > What are the relevant equations of motion in peophysics? What is the right description? Clustering of inertial particles + Bec, Gustavsson & Mehlig 2024


Particles in turbulent flows

Maxey and Riley (1983):

$$\mathbf{F}_{h} = \frac{4\pi}{3} \varrho_{f} a^{3} \frac{D}{Dt} \mathbf{u}(\mathbf{x}, t) - 6\pi \nu \varrho_{f} a \left[\mathbf{v} - \mathbf{u}(\mathbf{x}, t) \right] - \frac{4\pi}{3} \varrho_{f} a^{3} \mathbf{g}$$

$$- \frac{2\pi}{3} \varrho_{f} a^{3} \frac{d}{dt} \left[\mathbf{v} - \mathbf{u}(\mathbf{x}, t) \right] - 6\sqrt{\pi \nu} \varrho_{f} a^{2} \int_{0}^{t} \frac{ds}{\sqrt{t - s}} \frac{d}{ds} \left[\mathbf{v} - \mathbf{u}(\mathbf{x}, s) \right].$$

Coupling (see Elghobashi, 1994)



$$\int_{P} \frac{dv}{dt} = \frac{4\pi}{3} \rho_{t} a^{3} \frac{Du}{Dt} - 6\pi V \rho_{t} a \left(\underline{v} - \underline{u}\right) - \frac{4\pi}{3} \rho_{t} a^{3} \rho_{t}$$

$$-\frac{2\pi}{3} \rho_{t} a^{3} \frac{d}{dt} \left(\underline{v} - \underline{u}\right) - 6\sqrt{\pi} V \rho_{t} a^{2} \int_{V - S}^{t} \frac{ds}{ds} \frac{d}{\sqrt{t - s}} \frac{ds}{ds} \left(\underline{v} - \underline{u}\right)$$

$$\frac{drap}{dt}$$

$$\frac{dra$$

For very small a,
$$\rho_p \gg \rho_f$$
, $Re_p = \frac{\partial u - v}{\partial v} \ll 1$

$$\frac{\partial v}{\partial t} = \frac{1}{\zeta_p} \left(v - u \right) + g$$
with $\zeta_p = \frac{mp}{6\pi v \rho_f a}$ $m_p = \rho_p V_p$

$$1 \neq u = 0 \implies \frac{dv}{dt} = -\frac{1}{\zeta_p} v + g = 0$$

$$\implies v = \frac{\zeta_p g}{dt} = \frac{w}{\zeta_p}$$

$$1 \neq |\zeta_p| \ll |u| \implies \frac{dv}{dt} = -\frac{1}{\zeta_p} \left(v - u \right)$$

$$\dot{\sigma} = -\frac{1}{2}(\sigma - u)$$

we can decine à Stokes number

St =
$$\frac{Gp}{G}$$

But there isn't usually $G = G = (\frac{1}{2})^{1/2}$ an apreement on this!

Small eddies wou't effect the particle

Note

$$\underline{\sigma} = -\frac{1}{5}(\underline{\sigma} - \underline{u})$$
 is a low passband filter

Let's keep using MRG:
In a stratified flow:

$$\dot{y} = \frac{1}{\zeta_p} (\underline{u} - \underline{v}) - \frac{2N}{3} [N(z - z_0) - \zeta] \hat{z}$$
The first stratified flow:

$$+ \frac{Du}{Dt} + \sqrt{\frac{3}{\pi \zeta_{p}}} \int_{t-2}^{t} dz \frac{dz}{dz} (u-v)$$

Basset-Boussinesq

Added ______ buoyand (pravity)

mass _____ buoyand level (Zo)

Basset-Boursinesa

Basset-Boussinesquaditions this equation can be rewritten as


$$(\ddot{z}_p - \ddot{z}_f) + \frac{1}{5}(\dot{z}_p - \dot{z}_f) + \frac{2}{3}N^2(\ddot{z}_p - \ddot{z}_f) = F_{BB}^2$$

which is the equation for the forced damped oscillator.

For FBB = 0, we can pet + overdamped + critically damped + underdamped oscillations depending on Tp:

Using these solutions, we can bound $\frac{\|F_{BB} \cdot \hat{z}\|_{\infty}^{2}}{\|F_{SA} \cdot \hat{z}\|_{\infty}^{2}} \leq 16 N \zeta_{p}$


We can then define |Sb = NZp



Reartes & Mininni (2024)


In a rotating Flow:


$$\ddot{U} = \frac{1}{\zeta_{P}} \left(U - U \right) - \left(1 - \frac{f_{f}}{\rho_{P}} \right) p^{2} + \frac{3\rho_{f}}{\rho_{f} + 2\rho_{P}} \frac{Du}{Dt} - 2\Omega \times \left(U - \frac{3\rho_{f}}{\rho_{f} + 2\rho_{P}} \right) - \left(1 - \frac{3\rho_{f}}{\rho_{c} + 2\rho_{e}} \right) \Omega \times \Omega \times \left(X - X_{e} \right)$$




Biferale et al. (2016)

> Maxey's centrifuge Clusteriup Maxey 1987 Belkovsky-Felkovich-Fouxou 2001 Sweep-stick mechanism Goto & Vassilicos 2008 Other multiscale/12rge-scale Mechanisms (see Bec, Gustavssou, Mehlip 2024)





Obligado et al. (2014)



Karpińska et al. (2019)






Figure 3 Illustration of the evolution of a droplet size distribution during the onset of the collision-coalescence process. Figure adapted from Berry & Reinhardt (1974) and Lamb (2001), courtesy of D. Lamb, Penn State University.




Hunga Tonga-Hunga Ha'apai, January 15, 2022



Ring formation





From
$$\dot{y} = -\frac{1}{5}(y - u)$$

$$=$$
 $U = U - 72 + O(7^2)$

with
$$a = \frac{Du}{Dt} = \partial_t u + u \cdot \nabla u$$

Using that V.u=0

$$\Rightarrow \nabla \cdot \underline{\sigma} = - \overline{c}_p \operatorname{Tr}(A)$$

Regions with strain (Trs2>Tr22)
have V.0<0 => clusters
Regions dominated by rotation expell particles.

On the other hand, if Epis large, n= n-20 leads to particles selecting a=0, and these points are swept by the local fluid velocity u.

Preferential sampling

Maxey (1987): Maxey's centrifuge for St<1. Particles tend to be expelled from vortices.

Coleman and Vassilicos (2009): Sweep-stick mechanism. Particles accumulate at points of zero Lagrangian acceleration.

From Bec et al. (2014)

